

Microscopic colitis explained (MC for short)













Triggers

Medicines like: NSAIDs, PPIs Foods like: dairy, gluten, sweeteners



Treatments

Anti-diarrhoeals Bile salt binders Topical steroids (e.g. budesonide) There are other things you can do to help yourself





Don't smoke

Drink lots of water






Keep active


Talk things through






- MC is a type of inflammatory bowel disease with two main types: CC and LC¹
- In CC, there's a build-up of collagen in the lining of your gut (collagen is a threadlike protein that gives structure to your skin, bones and other body parts)^{2,3}
- In LC, there's a high number of lymphocytes in the lining of your gut (lymphocytes are a type of white blood cell that protects your body against disease)²
- We can tell them apart by looking at biopsies (small pieces of tissue taken during an endoscopy) under a microscope in the lab¹

In MC, the colon can't absorb all of the water from leftover waste, leading to watery diarrhoea





- Watery diarrhoea is the main symptom of MC, but we don't fully understand what causes it4
- The colon's main role is to absorb leftover water from the food you eat⁵
- When the colon is inflamed, it becomes less efficient at absorbing liquid from the waste, leading to watery stools⁶





Medicines like: NSAIDs, PPIs Foods like: dairy, gluten, sweeteners



Anti-diarrhoeals Bile salt binders **Topical steroids** (e.g. budesonide)

- Inflammation can go up and down, with periods of remission (when you feel well) and relapses (when you get symptoms)¹
- The first step to getting you into remission is taking away anything that might be triggering your MC4
 - NSAIDs (like aspirin and ibuprofen), PPIs (like omeprazole) and foods like dairy and sweeteners are common triggers4
- Most people will also need treatment that's tailored to their condition, like anti-diarrhoeals to control symptoms or steroids (like budesonide) to suppress the inflammation in your gut⁷

There are other things you can do to help yourself



Drink lots of water Don't smoke





Keep active

Talk things through

- Smoking can make it harder to get MC into remission, so it's best to avoid it7
- Regular exercise can help reduce fatigue and inflammation in people with IBD8
- Drink plenty of liquids to avoid dehydration when you have diarrhoea^{3,9}
- It's a good idea to talk things through - you can find lots of support through Crohn's & Colitis UK

References

Microscopic colitis (MC)

- 1. Miehlke S et al. Lancet Gastroenterol Hepatol 2019; 4: 305-14.
- 2. Münch A et al. J Crohns Colitis 2012; 6(9): 932-45.
- 3. National Institute of Diabetes and Digestive and Kidney Diseases. Microscopic Colitis. Available at: https://www.niddk.nih.gov/
- 4. Pardi DS. Am J Gastroenterol 2017; 112(1): 78-85.
- 5. Azzouz LL, Sharma S. Physiology, Large Intestine. Treasure Island, Florida: StatPearls Publishing, 2018.
- 6. Johns Hopkins Medicines. Collagenous & Lymphocytic Colitis: Introduction. Johns Hopkins University.
- 7. Boland K, Nguyen GC. Gastroenterol Hepatol 2017; 13(11): 671-7.
- 8. Bilski J et al. Pharmacol Rep 2016; 68(4): 827-36.
- 9. Brown AC et al. Expert Rev Gastroenterol Hepatol 2011; 5(3): 411-25.

Abbreviations

CC: collagenous colitis

IBD: inflammatory bowel disease

LC: lymphocytic colitis

MC: microscopic colitis

NSAID: non-steroidal anti-inflammatory drug

PPI: proton pump inhibitor

Useful links:

www.drfalk.co.uk www.crohnsandcolitis.org.uk

Adverse events should be reported. Reporting forms and information in Ireland can be found at www.hpra.ie and in the UK at www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. Adverse events should also be reported to Dr Falk Pharma UK Ltd. at PV@drfalkpharma.co.uk

Provided as a service to medicine by Dr Falk Pharma UK Ltd, Bourne End Business Park, Cores End Rd, Bourne End, SL8 5AS Registered in England No: 2307698

Date of preparation: September 2021

UI--2100111



